Test Bank for Calculus Early Transcendentals 9th Stewart

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

True / False
Indicate whether the statement is true or false.

___ 1. Determine whether the statement is true or false.

If $x_1 > x_2$ and f is a decreasing function, then $f(x_1) > f(x_2)$.

a. True

b. False

Multiple Choice
Indicate the answer choice that best completes the statement or answers the question.

___ 2. At the surface of the ocean, the water pressure is the same as the air pressure above the water, 13 lb/in^2. Below the surface, the water pressure increases by 4.34 lb/in^2 for every 10 ft of descent. At approximately what depth is the pressure 120 lb/in^2?

a. 247 ft

b. 276 ft

c. 30 ft

d. 56 ft

e. 359 ft


Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart


___ 3. Identify the graph of the following function.

$y = 5 - (0.2)^x$


a.




b.



c.



d.



Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 4. After alcohol is fully absorbed into the body, it is metabolized with a half-life of about 1.25 hours. Suppose you have had three alcoholic drinks and an hour later, at midnight, your blood alcohol concentration (BAC) is 0.6 mg/mL. Find an exponential decay model for your BAC t hours after midnight.

a. $\text{BAC}(t) = 0.6 \cdot (2)^{-\frac{4t}{5}}$

b. $\text{BAC}(t) = 1.25 \cdot (2)^{-\frac{t}{0.6}}$

c. $\text{BAC}(t) = 0.6 \cdot \left(\frac{1}{2}\right)^{1.25t}$

d. $\text{BAC}(t) = 1.25 \cdot \left(\frac{1}{2}\right)^{0.6t}$

e. None of the above

___ 5. The manager of a weekend flea market knows from past experience that if he charges x dollars for a rental space at the market, then the number y of spaces he can rent is given by the equation $y = 560 - 8x$. What does the x-intercept indicate?

a. There will be no spaces occupied when the rental cost is \$70.

b. There will be no spaces occupied when the rental cost is \$560.

c. There will be 560 spaces occupied when there is no rental cost.

d. There will be 8 spaces occupied when there is no rental cost.

e. The number of occupied spaces decreases by 5 for every \$1 increase in rent.

___ 6. Determine the range of the following function.

$g(x) = -2^{-x} - 9$

a. $(-\infty, -9)$

b. $(9, \infty)$

c. $\mathbb{R}$

d. $(-9, \infty)$

e. $(0, -\infty)$

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 7. Determine the domain and range of the function $f(x) = 3^x$.

a. Domain: $(-\infty, \infty)$, range: $(0, \infty)$
b. Domain: $(0, \infty)$, range: $(-\infty, \infty)$
c. Domain: $(0, \infty)$, range: $(0, \infty)$
d. Domain: $(-\infty, \infty)$, range: $(-\infty, \infty)$
e. Domain: $(-\infty, \infty)$, range: $(3, \infty)$
f. Domain: $(3, \infty)$, range: $(-\infty, \infty)$

___ 8. Starting with the graph of $y = e^x$, write the equation of the graph that results from shifting 7 units to the left and 6 units upward.

a. $y = e^{x+7} + 6$
b. $y = e^{x+6} + 7$
c. $y = e^{x+7} - 6$
d. $y = e^{x-7} + 6$
e. $y = 6e^{x-7}$

___ 9. Biologists have noticed that the chirping rate of crickets of a certain species is related to temperature, and the relationship appears to be very nearly linear. A cricket produces 108 chirps per minute at 70°F and 228 chirps per minute at 90°F. Find a linear equation that models the temperature T as a function of the number of chirps per minute N.

a. $T(N) = \frac{1}{6}N + \frac{312}{6}$
b. $T(N) = \frac{1}{6}N - \frac{312}{6}$
c. $T(N) = 6N - 648$
d. $N(T) = 6T - 312$
e. $N(T) = -6T + 312$

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 10. Find the function $f \circ g$ and its domain if $f(x) = \dfrac{x-1}{x}$ and $g(x) = \dfrac{x}{x+5}$.

a. $\dfrac{x-1}{x+5}, D = (-\infty, -5) \cup (-5, 0) \cup (0, \infty)$

b. $\dfrac{x-1}{x+5}, D = (-\infty, -5) \cup (-5, \infty)$

c. $-\dfrac{5}{x}, D = (-\infty, -5) \cup (-5, 0) \cup (0, \infty)$

d. $-\dfrac{5}{x}, D = (-\infty, 0) \cup (0, \infty)$

___ 11. Classify the following function:
$f(x) = x^6(9x^4 - 4)$

a. Polynomial function

b. Power function

c. Trigonometric function

d. Rational Function

e. Exponential function

___ 12. It makes sense that the larger the area of a region, the larger the number of species that inhabit the region. Many ecologists have modeled the species-area relation with a power function. In particular, the number of species S of bats living in caves in central Mexico has been related to the surface area A of the caves by the equation $S = 0.7A^{0.3}$. If you discover that nine species of bats live in a cave, estimate the area of the cave.

a. $4979\ \text{m}^2$

b. $1.35\ \text{m}^2$


c. $2125\ \text{m}^2$

d. $2.15\ \text{m}^2$

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 13. Decide what type of function you might choose as a model for the data displayed on the scatter plot.



a. exponential function
b. logarithmic function
c. polynomial
d. rational function
e. trigonometric function

___ 14. Find the function $f \circ g$ and its domain if $f(x) = \sqrt{x+6}$ and $g(x) = \sqrt{x-6}$.

a. $\sqrt{x^2 - 36}$
$D = [6, \infty)$
b. $\sqrt{x^2 - 36}$
$D = [-6, \infty)$
c. $\sqrt{x^2 + 36}$
$D = [-6, \infty)$
d. $\sqrt{x^2 + 36}$
$D = [6, \infty)$

___ 15. The half-life of bismuth-210, ^{210}Bi, is 5 days. If a sample has a mass of 600 mg, find the amount remaining after 4 weeks. Round your answer to one decimal place.

a. 12.4 mg
b. 29,102 mg
c. 0 mg
d. 82.8 mg
e. 6,000 mg


Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 16. If the recommended adult dosage for a drug is D (in mg), then to determine the appropriate dosage c for a child of age a, pharmacists use the equation $c = 0.04D(a + 1)$. Suppose the dosage for an adult is 900 mg. What is the slope of the graph of c and what does it represent?

a. 36 mg/yr; the rate of change of dosage with respect to age
b. 36 yr/mg; the rate of change of age with respect to dosage
c. 0.04 mg/yr; the rate of change of dosage with respect to age
d. 0.04 mg/yr; the rate of change of age with respect to dosage
e. 36 mg; the dosage for a newborn

___ 17. For the scatter plot below, what type of function might you see as a model for the data?



a. Trigonometric function
b. Rational Function
c. Exponential Function
d. Power Function
e. Polynomial Function

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 18. Find the domain of the following function.

$$f(x) = \frac{1 - e^{x^2}}{1 - e^{36 - x^2}}$$

a. $\mathbb{R}$


b. $x \neq 0$

c. $0 < x < 6$

d. $x \neq \pm 36$

e. $x \neq \pm 6$

___ 19. Let f be the function whose graph is given. Find the interval where the function is decreasing.



a. $(-\infty, 4)$

b. $(4, \infty)$

c. $[2, 6]$

d. $(-4, \infty)$

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 20. Use the Law of Exponents to rewrite and simplify the expression.

$$\frac{\sqrt{a\sqrt{b}}}{\sqrt[6]{ab}}$$

a. $\left(\frac{a}{b}\right)^{\frac{1}{3}}$

b. $\frac{a^{\frac{1}{3}}}{b^{\frac{1}{12}}}$

c. $\frac{a}{b}$

d. $\frac{a}{b^{\frac{1}{12}}}$

e. $a^{\frac{1}{3}} b^{\frac{1}{12}}$

___ 21. Identify the type of the following function.

$h(r) = 3.6^r$

a. power function
b. root function
c. polynomial
d. rational function
e. algebraic function
f. trigonometric function
g. exponential function
h. logarithmic function

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 22. Use the Law of Exponents to rewrite and simplify the expression.

$$\frac{(4z^8)^8}{2z^9}$$

a. $32{,}768z^{55}$

b. $65{,}536z^{55}$

c. $32{,}768z^{8}$

d. $2z^{55}$

e. $2z^{8}$

___ 23. An isotope of sodium, ^{24}Na, has a half-life of 15 hours. A sample of this isotope has mass 9 g. Estimate the time required for the mass to be reduced to 0.5 g.

a. 62.5 hours

b. 2.1 hours

c. 70.3 hours

d. 47.5 hours

e. 125.1 hours

___ 24. Find the domain of the function

$$f(x) = \frac{1}{\sqrt{3} - 2\cos x}$$

a. $x \neq \frac{\pi}{6} + 2\pi n$ and $x \neq \frac{11\pi}{6} + 2\pi n$, where n is any integer

b. $x \neq \frac{11\pi}{6}$ and $x \neq \frac{11\pi}{6}$

c. $x \neq \frac{5\pi}{6} + 2\pi n$ and $x \neq \frac{7\pi}{6} + 2\pi n$, where n is any integer

d. $x \neq \frac{5\pi}{6}$ and $x \neq \frac{7\pi}{6}$

e. $x \neq \frac{\pi}{4} + 2\pi n$ and $x \neq \frac{7\pi}{4} + 2\pi n$, where n is any integer

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 25. Find an equation for the family of linear functions such that $f(6) = 5$.

a. $f(x) = 5 + m(x - 6)$

b. $f(x) = m(x - 6) - 5$

c. $f(x) = 6 + m(x - 5)$

d. $f(x) = m(x + 5) - 5$

e. $f(x) = mx + 6m - 5$

___ 26. In the function $f(x) = 8x + d$, what must be the value d, if $f(3) = 1$?

a. $d = 25$

b. $d = 23$

c. $d = -23$

d. $d = -1$

e. $d = -25$


Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart


___ 27. Graph the function by hand, not by plotting points, but by starting with the graph of one of the standard functions and then applying the appropriate transformations.

$y = 3 + 2x - x^2$


a.




b.



c.



d.



Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 28. Use the Law of Exponents to rewrite and simplify the expression.

$$\frac{x^{4n} \cdot x^{5n+1}}{x^{n-4}}$$

a. x^{8n+5}

b. x^{8n+6}

c. x^{6n-9}

d. x^{3n+8}

e. x^{4n-7}

___ 29. Starting with the graph of $y = e^x$, find the equation of the graph that results from reflecting about the line $y = 5$.

a. $y = -e^x$

b. $y = -e^x + 10$

c. $y = -e^{-5x} + 10$

d. $y = e^{-x} + 10$

e. $y = -e^{x+10}$

___ 30. If $f(x) = x + 15$ and $h(x) = 4x + 85$, find a function g such that $g \circ f = h$.

a. $g(x) = 4x + 25$

b. $g(x) = 4x$

c. $g(x) = x - 25$


d. $g(x) = 4x - 25$

e. $g(x) = x + 25$

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 31. What is the equation of this graph?



a. $y = x^4$

b. $y = x^8$

c. $y = x^{10}$

d. $y = \sqrt[3]{x}$

e. $y = x^9$

___ 32. Suppose that the graph of f is given. Describe how the graph of the function $y = f(x - 3) - 3$ can be obtained from the graph of f.

a. Shift the graph 3 units to the left and 3 units down.

b. Shift the graph 3 units to the left and 3 units up.

c. Shift the graph 3 units to the right and 3 units up.

d. Shift the graph 3 units to the right and 3 units down.

e. None of these

___ 33. Find a, such that the function $f(x) = 4x + \sqrt{a - x^2}$ has the domain $[-3, 3]$.

a. $a = -9$

b. $a = \sqrt{3}$

c. $a = -\sqrt{3}$


d. $a = 9$

e. $a = 3$

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 34. Sandy wishes to have a rectangular garden in her backyard. She has 50 ft of fencing with which to enclose her garden. Letting x denote the width of the garden, find a function f in the variable x that gives the area of the garden.




a. $f(x) = 50x - x^2,\ 0 < x < 25$

b. $f(x) = 25x - x^2,\ 0 < x < 25$

c. $f(x) = 50x - x^2,\ 0 < x < 50$

d. $f(x) = 25x - x^2,\ 0 < x < 50$

___ 35. Which of the following is the equation for the function $g(x)$?



a. $g(x) = 4f(x)$

b. $g(x) = f(x) - 4$

c. $g(x) = -f(x + 4)$

d. $g(x) = \dfrac{f(x)}{4}$


e. $g(x) = -f(x) + 4$

Copyright Cengage Learning. Powered by Cognero.


Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 36. Sketch the graph of $y = 2 - \cos x$ over one period.


a.




b.



c.



d.




Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 37. A box with an open top is to be constructed from a rectangular piece of card board with dimensions $b = 6$ in. by $a = 25$ in. by cutting out equal squares of side x at each corner and then folding up the sides as in the figure.

Express the volume V of the box as a function of x.



a. $V(x) = x^3 - 62x^2 + 150x$

b. $V(x) = 4x^3 + 31x^2 + 196x$

c. $V(x) = 4x^3 + 62x^2 + 150x$

d. $V(x) = x^3 - 31x^2 + 196x$

e. $V(x) = 4x^3 - 62x^2 + 150x$

___ 38. Use the laws of logarithms to expand the expression.

$$\ln\left(\frac{x+1}{x-2}\right)^{\frac{1}{2}}$$

a. $\frac{1}{2}\ln[(x+1)(x-2)]$

b. $\frac{1}{2}\ln(x+1) - \frac{1}{2}\ln(x-2)$

c. $\frac{1}{2}\ln\left(x - \frac{1}{2}\right)$

d. $\frac{1}{2}\ln\left(\frac{x+1}{x-2}\right)$

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 39. The relationship between the Fahrenheit and Celsius temperature scales is given by the linear function.

$$F = \frac{9}{5}C + 32$$

What is the F-intercept and what does it represent?

a. $\frac{9}{5}$, Fahrenheit temperature corresponding to 0°C

b. $\frac{9}{5}$, Celsius temperature corresponding to 32°C

c. 32, Celsius temperature corresponding to 0°F

d. 0, Fahrenheit temperature corresponding to 32°C

e. 32, Fahrenheit temperature corresponding to 0°C

___ 40. Find an expression for the function $y = f(x)$ whose graph is the bottom half of the parabola $x + (10 - y)^2 = 0$.

a. $y = 10 - \sqrt{-x}$

b. $y = 100 - \sqrt{-x}$

c. $y = 10 + \sqrt{x}$

d. $y = 100 - x^2$


e. $y = 10 - x^2$

Copyright Cengage Learning. Powered by Cognero.


Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 41. Plot the graph of the function f in an appropriate viewing window.


$$f(x) = \frac{x^3}{x^3 + 1}$$




a.



b.



c.



d.

___ 42. Find $f^{-1}(a)$ for the function f and the real number a.

$$f(x) = x^3 + x - 2;\ a = -4$$

a. 0

b. –1

c. –2

d. 1

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 43. The monthly cost of driving a car depends on the number of miles driven. Julia found that in October it cost her \$300 to drive 400 mi and in July it cost her \$450 to drive 700 mi. Express the monthly cost C as a function of the distance driven d assuming that a linear relationship gives a suitable model.

a. $C = -100d + 0.5$
b. $C = 100d - 0.5$
c. $C = 0.5d + 100$
d. $C = 2d + 100$
e. $C = 0.5d - 100$

___ 44. Find the exact value of the given expression.

$$\cos\left(\cos^{-1}\left(-\frac{\sqrt{3}}{2}\right)\right)$$

a. 1
b. $-\frac{1}{2}$
c. 0
d. $-\frac{\sqrt{3}}{2}$

___ 45. If $f(x) = 5x^2 + 2$, find and simplify $\dfrac{f(1+h) - f(1)}{h}$, where $h \neq 0$.

a. $2 + 5h^2$
b. $5 + 10h$
c. $5h$
d. $10 + 5h$


___ 46. Find the inverse function of $f(x) = \dfrac{x + 1}{5x + 1}$.

a. $f^{-1}(x) = -(x + 1)(5x + 1)$
b. $f^{-1}(x) = -\dfrac{5x + 1}{x - 1}$
c. $f^{-1}(x) = -\dfrac{x - 1}{5x - 1}$
d. $f^{-1}(x) = \dfrac{x}{5x - 1}$
e. $f^{-1}(x) = (x + 1)(5x + 1)$

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 47. The graph of the function f is given. State the value of $f(2.7)$.



a. $f(2.7) = -40$


b. $f(2.7) = -10$

c. $f(2.7) = 40$

d. $f(2.7) = 0$

e. $f(2.7) = 10$

___ 48. The graph shown gives the weight of a certain person as a function of age. Find the age at which the person stopped an exercise program.



a. 20

b. 54

c. 38

d. 35

e. 70

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 49. If the point $(9, 7)$ is on the graph of an even function, what other point must also be on the graph?

a. $(-9, -7)$


b. $(9, -7)$

c. $(0, 0)$


d. $(-9, 7)$

e. None of these


___ 50. The graph of the function f follows. Choose the graph of $y = f(|x|)$.



a.




b.




Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

c.



d.



___ 51. Many physical quantities are connected by *inverse square laws*, that is, by power functions of the form:

$$f(x) = kx^{-2}$$

In particular, the illumination of an object by a light source is inversely proportional to the square of the distance from the source. Suppose that after dark you are in a room with just one lamp and you are trying to read a book. The light is too dim and so you move $\frac{2}{3}$ ds the distance to the lamp. How much brighter is the light?

a. $\frac{3}{2}$ times

b. $\frac{9}{4}$ times

c. $\frac{9}{2}$ times


d. $\frac{4}{9}$ times


e. 3 times

___ 52. The graph of the function f follows. Choose the graph of $y = -f(x) - 1$.


Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart





a.




b.


c.

d.

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart





___ 53. Solve the equation.

$3e^{x+2} = 2$

a. $x = \ln\frac{2}{3} - 2$

b. $x = \ln\frac{2}{3} + 2$

c. $x = \ln\frac{3}{2} - 2$

d. $x = \ln\frac{3}{2} + 2$

___ 54. Scientists have discovered that a linear relationship exists between the amount of flobberworm mucus secretions and the air temperature. When the temperature is 45°F, the flobberworms each secrete 8.5 grams of mucus a day; when the temperature is 75°F, they each secrete 11.5 grams of mucus a day. Find a function $M(t)$ that gives the amount of mucus secreted on a given day, where t is the temperature of that day in degrees Fahrenheit.

a. $M(t) = 0.1t + 8.5$

b. $M(t) = 10t + 8.5$

c. $M(t) = 10t + 4$

d. $M(t) = 0.1t + 4$

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 55. Starting with the graph of $y = e^x$, write the equation of the graph that results from shifting 6 units right.

a. $y = e^{x+6}$

b. $y = e^x - 6$

c. $y = e^{x-6}$

d. $y = e^x + 6$

e. $y = 6e^{x-6}$

___ 56. An open rectangular box with volume 4 m^3 has a square base. Express the surface area of the box as a function $S(x)$ of the length x of a side of the base.

a. $S(x) = x^2 + \dfrac{16}{x^2}$

b. $S(x) = x^2 + \dfrac{16}{x}$

c. $S(x) = 2x^2 + \dfrac{6}{x^2}$

d. $S(x) = x^2 + \dfrac{6}{x^2}$

e. $S(x) = 2x + \dfrac{6}{x}$

___ 57. Use the table to evaluate the expression $(f \circ g)(4)$.

x	1	2	3	4	5	6
$f(x)$	3	2	1	0	1	2
$g(x)$	6	5	2	3	4	6

a. 4

b. 1

c. 2

d. 3

e. 5

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 58. The graph of the function $f(x) = x^2 - 8x + 4$ has been shifted horizontally to the left by 9 units. Find the function for the transformed graph.


a. $g(x) = x^2 - 8x - 5$

b. $g(x) = x^2 - 8x + 13$

c. $g(x) = x^2 + 10x + 13$

d. $g(x) = x^2 - 26x + 157$

___ 59. Determine whether the function whose graph is given is even, odd, or neither.



a. Even

b. Neither

c. Odd


Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart


___ 60. Find the domain and sketch the graph of the function. What is its range?

$$f(x) = \begin{cases} -x + 3 & \text{if } x \geq 3 \\ x^2 - 2 & \text{if } x < 3 \end{cases}$$


a. ***D***: (–∞, ∞); ***R***: [–2, ∞)




b. ***D***: (–∞, ∞); ***R***: (–∞, 0]



c. ***D***: (–∞, ∞); ***R***: [0, ∞)



d. ***D***: (–∞, ∞); ***R***: (–∞, ∞)



___ 61. Determine whether the function is one-to-one.

$$f(x) = \sqrt{4 - x}$$

a. No

b. Yes

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 62. A rectangle has perimeter 14 m. Express the area of the rectangle as a function $A(l)$ of the length l of one of its sides.

a. $A(l) = 7l - l^2$

b. $A(l) = l - 7l^2$

c. $A(l) = 14l - l^2$

d. $A(l) = 14l + l^2$

e. $A(l) = 7l + l^2$

___ 63. If a bacteria population starts with 100 bacteria and doubles every three hours, then the number of bacteria after t hours is

$$n = f(t) = 100\left(2^{\frac{t}{3}}\right).$$

When will the population reach 45,000? Round the answer to the nearest tenth.

a. 26.4 hours

b. 16.4 hours

c. 31.4 hours

d. 21.4 hours

e. 36.4 hours

___ 64. Find the range of the function.

$y = 4 + \cos x$

a. $(-\infty, \infty)$

b. $(2, \infty)$

c. $[-1, 1]$

d. $(-1, 3)$

e. $[3, 5]$

___ 65. Classify the function as a Polynomial function, a Rational function, an algebraic function, or other.

$f(x) = x^9 + 4x^6 - 6$

a. Other

b. Algebraic

c. Polynomial

d. Rational


Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart


___ 66. Which of the following graphs is the graph of the function?

$f(x) = \sin|2x|$


Graph 1



Graph 2



Graph 3



a. Graph 2


b. Graph 1

c. Graph 3

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 67. The graphs of $f(x)$ and $g(x)$ are given. For what values of x is $f(x) = g(x)$?



a. 2, 3

b. 0

c. –1

d. –2, 5

e. –2, 10

___ 68. Find the domain of the function $f(x) = \dfrac{x}{7\sin x - 8}$

a. $(-\infty, \infty)$

b. $\left[\dfrac{8}{7}, \infty\right)$

c. $[-8, 7]$

d. $[-7, 8]$

___ 69. Find a formula for the inverse of the function.

$y = \ln(x + 8)$

a. $y = e^x - 8$

b. $y = e^x + 8$

c. $y = -8e^x$


d. $y = 8e^x$

e. $y = e^{x+8}$

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 70. Find the exponential function $y = Ca^x$ whose graph is given.



a. $f(x) = 2\left(\frac{1}{3}\right)^x$

b. $f(x) = 3\left(\frac{1}{3}\right)^x$

c. $f(x) = \left(\frac{3}{2}\right)$

d. $f(x) = \frac{1}{3^x}$

e. $f(x) = \left(\frac{3}{4}\right)^x$

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 71. Use the laws of logarithms to write the expression as the logarithm of a single quantity.

$$4 \ln 5 - \frac{3}{4}\ln(x + 3)$$

a. $\ln \dfrac{625}{(x + 3)^{3/4}}$

b. $\dfrac{4 \ln 5}{\frac{3}{4}\ln(x + 3)}$

c. $\ln 5^4 - \ln(x + 3)^{3/4}$

d. $\dfrac{3 \ln 625}{4(x + 3)}$

___ 72. Which of the following functions is neither even nor odd?

a. $f(x) = \dfrac{2x^2}{x^4 + 1}$

b. $f(x) = 8x^3 + 10x^2 + 7$

c. $f(x) = x^3 - 3x$

___ 73. Find the domain of the function.

$$f(x) = \frac{7x + 5}{x^2}$$

a. $(-\infty, 0)$

b. $\left(-\infty, -\dfrac{5}{7}\right) \cup \left(-\dfrac{5}{7}, \infty\right)$

c. $\left(-\infty, \dfrac{5}{7}\right) \cup \left(\dfrac{5}{7}, \infty\right)$

d. $(-\infty, 0) \cup (0, \infty)$

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 74. Find the exact value of the expression.

$\log_5 100 + \log_5 50 - 3\log_5 2$

a. 7
b. 8
c. 6
d. 4
e. 5

___ 75. Under ideal conditions a certain bacteria population is known to double every three hours. Suppose that there are initially 20 bacteria. What is the size of the population after 15 hours?

a. 1,280 bacteria
b. 80 bacteria
c. 640 bacteria
d. 160 bacteria
e. 320 bacteria

___ 76. If $f(x) = x^2 - x + 8$, evaluate the difference quotient $\dfrac{f(a+h) - f(a)}{h}$.

a. $2a + h - 8$
b. $2a - 8$
c. $2a - h - 8$
d. h
e. none of these

___ 77. The table gives the population of the United States, in millions, for the years 1900 - 2000. Use a graphing calculator with exponential regression capability to model the U.S. population since 1900. Use the model to estimate the population in 1965 and to predict the population in the year 2025.

Year	1900	1910	1920	1930	1940	1950	1960	1970	1980	1990	2000
Population (millions)	76	92	106	123	131	150	179	203	227	250	281

a. 185 million, 400 million
b. 190 million, 405 million
c. 195 million, 410 million
d. 180 million, 395 million
e. 175 million, 390 million

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 78. Simplify the expression.

$e^{3\ln 4}$

a. 7
b. 12
c. 64
d. 81

___ 79. Find the exact value of the given expression.

$\tan^{-1} 1$

a. 2π
b. $\frac{\pi}{4}$
c. $\frac{4}{\pi}$
d. 4π

Copyright Cengage Learning. Powered by Cognero.


Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 80. Find the inverse of f. Then sketch the graphs of f and f^{-1} on the same set of axes.

$f(x) = \sqrt{16 - x^2}, x \geq 0$


a. $f^{-1}(x) = -\sqrt{16 - x^2}, x \geq 0$

$f = f^{-1}$




b. $f^{-1}(x) = \sqrt{4 + x^2}, x \geq 0$

$f(blue), f^{-1}(red)$




c. $f^{-1}(x) = -\sqrt{4 + x^2}, x \geq 0$

$f(blue), f^{-1}(red)$



d. $f^{-1}(x) = \sqrt{16 - x^2}, x \geq 0$

$f = f^{-1}$



___ 81. Suppose that the graph of $y = \log_2 x$ is drawn on a coordinate grid where the unit of measurement is an inch. How many miles to the right of the origin do we have to move before the height of the curve reaches 3 ft? Rounded to the nearest mile.

a. 5.2 mi

b. 1,084,587.7 mi

c. 103 mi

d. 13,015,052.4 mi

e. –97 mi

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 82. Determine whether f is even, odd, or neither.

$$f(x) = \frac{8x^2}{x^4 + 5}$$

a. neither
b. odd
c. even

___ 83. Find the domain.

$$g(u) = \sqrt{u} - \sqrt{3 - u}$$

a. $(0, 3)$
b. $(-\infty, 0]$
c. $(-3, \infty]$
d. $[0, 3]$
e. $[0, \infty)$

___ 84. What is $\sqrt[10]{x}$, given that $H = h \circ f \circ g$ and $H(x) = \sqrt[10]{\sqrt{x} - 3}$?

a. $g(x)$
b. $f(x)$
c. $h(x)$

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

___ 85. When a camera flash goes off, the batteries immediately begin to recharge the flash's capacitor, which stores electric charge given by

$$Q(t) = Q_0\left(1 - e^{-t/a}\right)$$

(The maximum charge capacity Q_0 is and t is measured in seconds.) How long does it take to recharge the capacitor to 90% of capacity if a = 4?

a. $-4\ln\left(\frac{9}{10}\right)$ seconds

b. $-4\ln\left(\frac{1}{10}\right)$ seconds

c. $-4\ln(10)$ seconds

d. $-4\ln(1)$ seconds

e. $-\ln\left(\frac{4}{9}\right)$ seconds

Numeric Response

Enter the appropriate value to answer the question or solve the problem.

86. Simplify the expression.

$$\sin(2\cos^{-1}4x)$$

87. A spherical balloon with radius r inches has volume $4\frac{\pi r^3}{3}$.

Find a function that represents the amount of air required to inflate the balloon from a radius of r inches to a radius of $r + 2$ inches.

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

88. If $f(x) = 3x + \ln x$, find $f^{-1}(3)$. Round to the nearest whole.

89. The relationship between the Fahrenheit and Celsius temperature scales is given by the linear function.

$$F = \frac{9}{5}C + 32$$

Complete the table and find the slope. Round your answers to the nearest whole number.

°C	**°F**
15	
–15	
slope	

90. Express the function in the form of $f \circ g \circ h$.

$$H(x) = 3 - 5^{x^3}$$

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

91. Find the domain of the function.

$f(x) = \sqrt{25 - x^2}$

92. Fill in the blanks.

Let $f(x) = 3 + x^2 + \tan\left(\frac{\pi x}{2}\right)$, where $-1 < x < 1$.

$f^{-1}(f(3))$_____ $f(f^{-1}(5))$_____

93. A stone is dropped into a lake, creating a circular ripple that travels outward at a speed of 49 cm/s. Express the radius r of this circle as a function of the time t (in seconds) and find $A \circ r$, if A is the area of this circle as a function of the radius.

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

94. It makes sense that the larger the area of a region, the larger the number of species that inhabit the region. Many ecologists have modeled the species-area relation with a power function and, in particular, the number of species S of bats living in caves in central Mexico has been related to the surface area A measured in m^2 of the caves by the equation

$$S = 0.7A^{0.3}$$

(a) The cave called mission impossible near puebla, mexico, has suface area of $A = 80m^2$.

How many species of bats would expect to find in that cave?

(b) If you discover that 2 species of bats live in cave estimate the area of the cave.

95. Express the function in the form of $f \circ g$.

$$v(t) = \tan(t^2)\sin(t^2)$$

96. A spherical balloon with radius r inches has volume $\frac{4}{3}\pi r^3$.

Find a function that represents the amount of air required to inflate the balloon from a radius of r inches to a radius of $r + 7$ inches.

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

97. Solve each equation for x.

(a) $\ln x = 3$ (b) $e^{e^x} = 6$

98. Find the range of the function.

$h(x) = \sqrt{36 - x^2}$

99. Find the exact value of the expression.

$\cos\left(\arcsin\frac{1}{2}\right)$

100. Find the domain of the function.

$f(x) = \frac{3}{1 - e^x}$


Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

101. The graphs of $f(x)$ and $g(x)$ are given.

a) For what values of x is $f(x) = g(x)$?

b) Find the values of $f(-1)$ and $g(12)$.



102. Find the domain of the function.

$$f(x) = \frac{7}{3x - 1}$$

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

103. Determine whether f is even, odd, or neither.


$$f(x) = \frac{2x^2}{x^4 + 1}$$

104. Find the domain, range, and x-intercept(s) of the function.

$$f(x) = \ln(x - 4) - 1$$

Subjective Short Answer


105. Use the vertical line test to determine whether the curve is the graph of a function of x.



Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

106. The following figure shows a portion of the graph of a function f defined on the interval $[-1, 1]$. Sketch the complete graph of f if it is known f is odd.




107. Find the function g such that $h(x) = (g \circ f)(x)$.

$h(x) = \sin^5 x$ and $f(x) = \sin x$

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart


108. By cutting away identical squares from each corner of a rectangular piece of cardboard and folding up the resulting flaps, an open box can be made. If the cardboard is 18 in. long and 9 in. wide and the square cutaways have dimensions of x in. by x in., find a function that gives the volume of the resulting box.



Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

109. Refer to the graph of the function f in the following figure.



a. Find $f(0)$.
b. Find the value of x for which (i) $f(x) = 1$ and (ii) $f(x) = 0$.
c. Find the domain and range of f.

110. Sketch the graph of f.

$f(x) = \ln(x + 5) - 3$


111. Sketch the graph of the function.

$f(x) = |1 - 7x|$

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

112. Determine whether f is one-to-one.



113. Find the inverse of f. Then use a graphing utility to plot the graphs of f and f^{-1} on the same set of axes.

$$f(x) = \frac{1 - e^{6x}}{1 + e^{6x}}$$

114. If $f(x) = \begin{cases} x^2 + 7 & \text{if } x \le 0 \\ \sqrt{x} & \text{if } x > 0 \end{cases}$

find $f(-3)$, $f(0)$, and $f(4)$.

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

115. Find $f \circ g \circ h$ if

$f(x) = \sqrt{x}$, $g(x) = 3x + 1$, and $h(x) = x^2 - 1$

116. Determine whether the function is even, odd, or neither.

$f(x) = 4x^2 + 8x + 1$


117. Let $f(x) = x^2 - 18x + 81$ and $g(x) = \sqrt{x + 1}$. Find $(f \circ f)(12)$.

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

118. The graph of f is given. Sketch the graph of f^{-1} on the same set of axes.

$y = f(x)$




119. Find the inverse of f. Then sketch the graphs of f and f^{-1} on the same set of axes.

$$f(x) = \sin^{-1}\left(\frac{x}{5}\right), \quad -5 \le x \le 5$$

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

120. The graph of a function g is given. On what interval(s) is g increasing?



Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

Answer Key

1. False

2. a

3. a

4. a

5. a

6. a

7. a

8. a

9. a

10. c

11. a

12. a

13. a

14. a

15. a

16. a

17. a

18. e

19. a

20. e

21. g

22. a

23. a

24. a

25. a

26. c

27. c

28. a

29. b

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

30. a

31. e

32. d

33. d

34. b

35. d

36. c

37. e

38. b

39. e

40. a

41. c

42. b

43. c

44. d

45. d

46. c

47. c

48. d

49. d

50. d

51. b

52. a

53. a

54. d

55. c

56. b

57. b

58. c

59. c

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

60. d

61. b

62. a

63. a

64. e

65. c

66. a

67. e

68. a

69. a

70. a

71. a

72. b

73. d

74. d

75. c

76. e

77. a

78. c

79. b

80. d

81. b

82. c

83. d

84. c

85. b

86. $8x\sqrt{1 - 16x^2}$

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

87. $\frac{8}{3}\pi(6r^2+12r+8)$

88. $f^{-1}(3) = 1$

89. (15, 59), (–15, 5); slope = 2

90. $h(x)=x^3, g(x)=5^x, f(x)=3-x$

91. [–5, 5]

92. 3, 5

93. $r(t)=49t,\ 2401\,\pi\,t^2$

94. a) 3 species
b) 33 m^2

95. $f(t)=\tan(t)\sin(t)$
$g(t)=t^2$

96. $\frac{28}{3}\pi(3r^2+21r+49)$

97. $x = e^3, x = \ln(\ln 6)$

98. $0 \le h(x) \le 6$

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

99. $\frac{\sqrt{3}}{2}$

100. $(-\infty, 0) \cup (0, \infty)$


101. a) 2, 6 b) $f(-1) = 12, g(12) = -10$

102. $\left\{x \middle| x \neq \frac{1}{3}\right\}$

103. even

104. Domain: $(4, \infty)$ Range: $(-\infty, \infty)$ x-intercept: $x = e + 4$

105. Yes



106.

107. $g(x) = x^5$

108. $V = 4x^3 - 54x^2 + 162x$


Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart


109. **a.** 0

b. (**i**) 1 (**ii**) 0, 2

c. ***D***: [0, 3], ***R***: [–3, 1]



110.




111.

112. No

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

113. $f^{-1}(x) = \frac{1}{6}\ln\left(\frac{1-x}{1+x}\right)$




114. $f(-3) = 16,\ f(0) = 7,\ f(4) = 2.$

115. $\sqrt{3x^2 - 2}$

116. Neither

117. 0


118. $y = f(x)$ - blue, $y = f^{-1}(x)$ -red



Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart

119. $f^{-1}(x) = 5\sin x,\ -\frac{\pi}{2} \le x \le \frac{\pi}{2}$



120. $[-6, -4] \cup (-2, 2]$

Copyright Cengage Learning. Powered by Cognero.

Access Complete Test Bank for Calculus Early Transcendentals 9th Stewart